

# Catalog of Fishes Queries May 2016

Dennis Polack Fishwisepro

*epuluensis*, *Brycinus* Decru [E.], **Vreven [E.]**, Sadio [O.] & Snoeks [J.] 2016:55, Figs. 4, 5 [Ichthyological Exploration of Freshwaters v. 27 (no. 1); ref. [34370](#)] Edoro River, upstream of the bridge, near the research camp at Edoro-Afarama, Epulu River, Ituri River basin, Democratic Republic of Congo, 1°33'01"N, 28°30'41"E. Holotype: MRAC A9-029-P-0612. Paratypes: MRAC. Plus additional non-type material. •Valid as *Brycinus epuluensis* Decru, Vreven, Sadio & Snoeks 2016. **Current status:** Valid as *Brycinus epuluensis* Decru, Vreven, Sadio & Snoeks 2016. Alestidae. Distribution: Epulu River, Ituri River basin, Democratic Republic of Congo. Habitat: freshwater.

**Usually Vreven [E. J.]**,

:

*guercifensis*, *Luciobarbus* **Doadrio [I.]**, **Perea [S.] & Yahyaoui [A.]** **Doadrio [I.]**, **Casal-López [M.]**, **Perea [S.] & Yahyaoui [A.]** 2016:8, Fig. 7 [Graellsia v. 72 (no. 1, e039); ref. [34185](#)]

Moulouya River, Moulouya Basin, Ghafoula, Mediterranean slope in Morocco, 34°08'43.2"N, 3°23'18.5"W, elevation 378 meters. Holotype: MNCN 290835. Paratypes: MNCN. •Valid as *Luciobarbus guercifensis* Doadrio, Perea & Yahyaoui 2016. **Current status:** Valid as *Luciobarbus guercifensis* Doadrio, Perea & Yahyaoui 2016. Cyprinidae: Cyprininae. Distribution: Moulouya Basin, Morocco. Habitat: freshwater. **Doadrio [I.]**,

**Casal-López [M.]**, **Perea [S.] & Yahyaoui [A.]**

:

*multisquamatus*, *Marcusenius* **Kramer [B.]** & Wink [M.] 2013:13 [2339], Fig. 4e [Journal of Natural History v. 47 (nos. 35-36); ref. 32824] Cunene River: Epupa Falls, Hot Springs Campsite, estimated 300 meters upstream from the Falls, 17°00'07"S, 13°14'57"E, Namibia, elevation about 600 meters. Holotype: SAIAB 78781. Paratypes: SAIAB 78780 (2), 78792 (1); ZSM 38526 (2), 38527 (2). Plus non-type specimens. •Valid as *Marcusenius multisquamatus* Kramer & Wink 2013 -- (Maake et al. 2014:466 [ref. 33223], Kramer et al. 2016:8 [ref. 34331]). Current status: Valid as *Marcusenius multisquamatus* Kramer & Wink 2013. Mormyridae. Distribution: Cunene River just above Epupa Falls, Angola/Namibia border. Habitat: freshwater.

**This Kramer normally has the initials [L. B.]**

:

*altisambesi*, *Marcusenius* **Kramer [B.]**, Skelton [P. H.], van der Bank [F. H.] & Wink [M.] 2007:681, Fig. 1K [Journal of Natural History v. 41 (9-12); ref. 29168] Caprivi Strip, Upper Zambezi River at Lisikili, Namibia. Holotype: SAIAB 79135. Paratypes: SAIAB 79136 (6), 79137 (3); ZSM 35082 (2), 35085 (2), 35086 (5), 35097 (1). Type catalog: Neumann 2011:270 [ref. 31652]. •Valid as *Marcusenius altisambesi* Kramer, Skelton, van der Bank & Wink 2007 -- (Kramer & van der Bank 2011:2670 [ref. 31546], Marshall 2011:52 [ref. 31755], Kramer & Wink 2013:9 [2335] [ref. 32824], Maake et al. 2014:466 [ref. 33223], Kramer et al. 2016:6 [ref. 34331]). Current status: Valid as *Marcusenius altisambesi* Kramer, Skelton, van der Bank & Wink 2007. Mormyridae. Distribution: Upper Zambezi and Okavango River systems, Africa. Habitat: freshwater.

**This Kramer normally has the initials [L. B.]**

:

devosi, Marcusenius **Kramer [B.]**, Skelton [P. H.], van der Bank [F. H.] & Wink [M.] 2007:686, Fig. 1G [Journal of Natural History v. 41 (9-12); ref. 29168] Lower Tana River near village Wenje, Kenya, 1°52'38.1"S, 40°08'22.5"E. Holotype: SAIAB 79138. Paratypes: SAIAB 79139 (14); ZSM 35091 (1), 35092 (1), 35093 (4), 35094 (7). Type catalog: Neumann 2011:271 [ref. 31652]. •Valid as Marcusenius devosi Kramer, Skelton, van der Bank & Wink 2007 -- (Kramer & Wink 2013:8 [2344] [ref. 32824], Maake et al. 2014:466 [ref. 33223], Kramer et al. 2016:7 [ref. 34331]). Current status: Valid as Marcusenius devosi Kramer, Skelton, van der Bank & Wink 2007. Mormyridae. Distribution: Tana River, Kenya. Habitat: freshwater.

**This Kramer normally has the initials [L. B.]**

**auriferus, Prorivulus** Costa [W. J. E. M.], Lima [S. M. Q.] & **Suzart [R.]** in Costa 2004:3, Figs. 1-2 [Zootaxa No. 642; ref. 27874] Valença, coastal stream, (24L 0503477 UTM 8505050), Estado da Bahia, Brazil. Holotype: UFRJ 5932. Paratypes: UFRJ 5933 (4), 5934 (3, c&s). •Valid as *Prorivulus auriferus* Costa, Lima & Suzart 2004. **Current status:** Valid as *Prorivulus auriferus* Costa, Lima & Suzart 2004. Rivulidae. Distribution: Brazil. Habitat: freshwater.

**CoF has several variations for author "Suzart". I have standardized on "dos Reis Suzart [R.]"**

:

**macrolepis, Mugil** Bleeker [P.] 1852:422 [Natuurkundig Tijdschrift voor Nederlandsch Indië v. 3 (no. 3); ref. 16827] Pamangkat rivers, Sampit (marine), Borneo, Indonesia. Syntypes: (4). Objectively invalid; preoccupied by *Mugil macrolepis* Smith 1846. •Synonym of *Liza parmatus* (Cantor 1859) -- (Thomson 1997:528 [ref. 23428] dated 1850, Kottelat 2013:277 [ref. 32989]). •Synonym of *Planiliza parmatus* (Cantor 1859). **Current status:** Synonym of **Planiliza** *parmatus* (Cantor 1849). Mugilidae.

**Paramugil**

:

**eurystomus, a** var. Keilhack [L.] 1908:166 [Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin 1908 (no. 7); ref. 14324] Langenburg, Lake Malawi [Lake Nyasa], southeastern Africa. Lectotype: ZMB 18175. Paralectotypes: ZMB 18175 (2). Lectotype apparently designated first by Keilhack 1910:103 [ref. 32109]. Also appeared as new in Keilhack 1910:102 [ref. 32109]. •Valid as *Barbus eurystomus* Keilhack 1908 -- (Lévêque & Daget 1984:243 [ref. 6186]). •Synonym of *Labeobarbus brevicauda* (Keilhack 1908) -- (Seegers 1995:31 [ref. 33299]). •Synonym of *Labeobarbus johnstonii* (Boulenger 1907) -- (Vreven et al. 2016:30, 61 [ref. 34168]). **Current status:** Synonym of *Labeobarbus johnstonii* (Boulenger 1907). Cyprinidae: Cyprininae. Habitat: freshwater.

**eurystomus, Barbus (Probably Barbus)**

: